

REFERENCES

- [1] Wielage RC, Bansal M, Andrews JS, Wohlreich MM, et al. The cost-effectiveness of duloxetine in chronic low back pain: a US private payer perspective. *Value Health* 2013;16:334–44.
- [2] Birbara CA, Puopolo AD, Munoz DR, et al. Treatment of chronic low back pain with etoricoxib, a new cyclo-oxygenase-2 selective inhibitor: improvement in pain and disability—a randomized, placebo-controlled, 3-month trial. *J Pain* 2003;4:307–15.
- [3] Palla RM, Seger W, Adler JL, et al. Etoricoxib reduced pain and disability and improved quality of life in patients with chronic low back pain: a 3 month, randomized, controlled trial. *Scand J Rheumatol* 2004;33:257–66.
- [4] Kelly J, Cook S, Kaufman D, et al. Prevalence and characteristics of opioid use in the US adult population. *Pain* 2008;138:507–13.
- [5] Williams R, Bosnic N, Sweeney C, et al. Prevalence of opioid dispensings and concurrent gastrointestinal medications in Quebec. *Pain Res Manage* 2008;13:395–400.
- [6] Dart RC, Cicero TJ, Surratt HL, et al. Assessment of the abuse of tapentadol immediate release: the first 24 months. *J Opioid Manag* 2012;8:395–402.

ERRATA

In: Vemer P, Krabbe PFM, Feenstra TL, van Voorn GAK, et al. Improving model validation in HTA: comments on the guidelines of the ISPOR-SMDM Modeling Good Research Practices Task Force. *Value Health*. 2013;16:1106–7.

The correct authors listing should be as follows: Pepijn Vemer, MSc, George A.K. van Voorn, PhD, Isaac Corro Ramos, PhD, Paul F.M. Krabbe, PhD, Maiwenn J. Al, PhD, Talitha L. Feenstra, PhD

References 4 and 5 should appear as follows:

[4] Peters B, Smith J, Medeiros D, Rohrer M, eds. How to build valid and credible simulation models. In: 2001 Winter Simulation Conference, Arlington, VA, December 9–12, 2001.

[5] Ingalls R, Rossetti M, Smith J, Peters B, eds. Validation and verification of simulation models. In: 2004 Winter Simulation Conference, Washington, D.C., December 5–8. Piscataway, NJ: IEEE Press, 2004.

In: Abstract PMH19, “Association between cognitive function and 3 month healthcare costs among patients initiating an antidepressant for depressive disorder in an ambulatory care setting” by Kurlander JL, MS¹; Walker V, Essoi B, Samp JC, Yang J, Akhras KS (Value Health 2013;16:A544). This poster was presented at the ISPOR 16th Annual European Congress in Dublin, Ireland on November 5th, 2013.

The correct full text of the abstract appears below.

PMH19

ASSOCIATION BETWEEN COGNITIVE FUNCTION AND 3 MONTH HEALTHCARE COSTS AMONG PATIENTS INITIATING AN ANTIDEPRESSANT FOR DEPRESSIVE DISORDER IN AN AMBULATORY CARE SETTING

Kurlander JL, MS¹; Walker V McS¹; Essoi B, MPH¹; Samp JC, PharmD, MS²; Yang, J, PhD²; Akhras KS, PharmD²

¹OptumInsight, MN002-0258, 12125 Technology Drive, Eden Prairie, MN 55344, ²Takeda Pharmaceuticals Inc., One Takeda Parkway, Deerfield, IL 60015

Objectives: Depression is associated with reduced cognitive function and significant healthcare costs; however, the extent to which these two are related remains unclear. This study compared follow-up healthcare costs for major depressive disorder patients with and without cognitive dysfunction after antidepressant (AD) initiation. **Methods:** A large US health plan affiliated with OptumInsight was used to identify depressed patients with a newly prescribed AD who could be surveyed to assess cognitive function. Patients with neurological diseases associated with cognitive dysfunction were excluded. Patients were mailed a survey invitation and consent form. Patients maintained eligibility by confirming a depressive diagnosis and no excluding diagnoses. Consenting, eligible patients were interviewed by telephone and completed 4 cognitive function tests. Patients were classified as “cognitive normal (CN)” or “cognitive dysfunction (CD)” based on test scores relative to normative data. All-cause healthcare costs in the 3 months post-AD initiation were calculated from pharmacy and medical claims. T-tests compared 3-month costs of CN versus CD. Gamma models with log link compared healthcare costs between CD and CN patients, adjusting for race, sex, age, education, employment, depression severity, and comorbidities. **Results:** 13,537 patients were invited to participate in the study and 564 patients maintained eligibility and completed the study. Patients were mostly female (80%), mean age was 41 years, 98% had a high school degree or higher, and 84% were employed. A total of 45% (n=255) met criteria for CD. Mean healthcare costs were \$3,053 for all patients. Costs were \$3,948 for the CD group compared to \$2,312 for the CN (p = 0.113). In the gamma models with costs as the outcome, CD patients had costs 1.46 times higher than CN patients (95% CI: 1.12, 1.92)(p=0.0059). **Conclusions:** In this study population, healthcare costs were significantly higher in patients with cognitive dysfunction compared to those without cognitive dysfunction.

Commercial Support Disclosure: The study was jointly funded by Takeda Pharmaceutical Company and H. Lundbeck A/S.